

```

> restart
> Sist := diff(x[1](t), t) = x[1](t) + 2·x[2](t) + 5·exp(3·t), diff(x[2](t), t) = 4·x[1](t) + 3
    ·x[2](t) - 7·exp(2·t) : Sist[1]; Sist[2]
    
$$\frac{d}{dt} x_1(t) = x_1(t) + 2 x_2(t) + 5 e^{3t}$$

    
$$\frac{d}{dt} x_2(t) = 4 x_1(t) + 3 x_2(t) - 7 e^{2t} \quad (1)$$

> Cond := x[1](0) = 2, x[2](0) = -2 : Cond[1]; Cond[2]
    
$$x_1(0) = 2$$

    
$$x_2(0) = -2 \quad (2)$$

> BB := 5·exp(3·t), -7·exp(2·t) : BB[1]; BB[2]
    
$$5 e^{3t}$$

    
$$-7 e^{2t} \quad (3)$$

> with(LinearAlgebra) :
> with(inttrans) :
> with(linalg) :
> AA := Matrix([ [1, 2], [4, 3] ])
    
$$AA := \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix} \quad (4)$$

> II := Matrix([ [1, 0], [0, 1] ])
    
$$II := \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad (5)$$

> MatUno := evalm(s·II - AA)
    
$$MatUno := \begin{bmatrix} s-1 & -2 \\ -4 & s-3 \end{bmatrix} \quad (6)$$

> InvMatUno := inverse(MatUno)
    
$$InvMatUno := \begin{bmatrix} \frac{s-3}{s^2-4s-5} & \frac{2}{s^2-4s-5} \\ \frac{4}{s^2-4s-5} & \frac{s-1}{s^2-4s-5} \end{bmatrix} \quad (7)$$

> MatExp := map(expand, (map(convert, map(invlaplace, InvMatUno, s, t), exp)))
    
$$MatExp := \begin{bmatrix} \frac{1}{3} (e^t)^5 + \frac{2}{3} e^t & \frac{1}{3} (e^t)^5 - \frac{1}{3} e^t \\ \frac{2}{3} (e^t)^5 - \frac{2}{3} e^t & \frac{2}{3} (e^t)^5 + \frac{1}{3} e^t \end{bmatrix} \quad (8)$$

> MatExpDos := exponential(AA, t)
    
$$(9)$$


```

$$MatExpDos := \begin{bmatrix} \frac{1}{3} e^{5t} + \frac{2}{3} e^{-t} & \frac{1}{3} e^{5t} - \frac{1}{3} e^{-t} \\ \frac{2}{3} e^{5t} - \frac{2}{3} e^{-t} & \frac{2}{3} e^{5t} + \frac{1}{3} e^{-t} \end{bmatrix} \quad (9)$$

>  $MatExpTres := MatrixExponential(AA, t)$

$$MatExpTres := \begin{bmatrix} \frac{1}{3} e^{5t} + \frac{2}{3} e^{-t} & \frac{1}{3} e^{5t} - \frac{1}{3} e^{-t} \\ \frac{2}{3} e^{5t} - \frac{2}{3} e^{-t} & \frac{2}{3} e^{5t} + \frac{1}{3} e^{-t} \end{bmatrix} \quad (10)$$

>  $Identidad := map(rcurry(eval, t='0'), MatExpDos)$

$$Identidad := \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad (11)$$

>  $DerMatExp := map(diff, MatExpDos, t)$

$$DerMatExp := \begin{bmatrix} \frac{5}{3} e^{5t} - \frac{2}{3} e^{-t} & \frac{5}{3} e^{5t} + \frac{1}{3} e^{-t} \\ \frac{10}{3} e^{5t} + \frac{2}{3} e^{-t} & \frac{10}{3} e^{5t} - \frac{1}{3} e^{-t} \end{bmatrix} \quad (12)$$

>  $Comprobacion := evalm(DerMatExp - evalm(AA &* MatExpDos))$

$$Comprobacion := \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \quad (13)$$

>  $IdentidadDos := simplify(evalm(map(rcurry(eval, t=-t'), MatExpDos) &* MatExpDos))$

$$IdentidadDos := \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad (14)$$

>  $evalm(MatExpDos)$

$$\begin{bmatrix} \frac{1}{3} e^{5t} + \frac{2}{3} e^{-t} & \frac{1}{3} e^{5t} - \frac{1}{3} e^{-t} \\ \frac{2}{3} e^{5t} - \frac{2}{3} e^{-t} & \frac{2}{3} e^{5t} + \frac{1}{3} e^{-t} \end{bmatrix} \quad (15)$$

>  $Xcero := array([ [2], [-2] ])$

$$Xcero := \begin{bmatrix} 2 \\ -2 \end{bmatrix} \quad (16)$$

>  $MatExpHom := evalm(MatExpDos &* Xcero)$

$$MatExpHom := \begin{bmatrix} 2 e^{-t} \\ -2 e^{-t} \end{bmatrix} \quad (17)$$

>  $BBtau := map(rcurry(eval, t='tau'), array([ [BB[1]], [BB[2]]]))$

(18)

$$BBtau := \begin{bmatrix} 5 e^{3\tau} \\ -7 e^{2\tau} \end{bmatrix} \quad (18)$$

>  $MatExpTau := map(rcurry(eval, t=t-tau'), MatExpDos)$

$$MatExpTau := \begin{bmatrix} \frac{1}{3} e^{5t-5\tau} + \frac{2}{3} e^{-t+\tau} & \frac{1}{3} e^{5t-5\tau} - \frac{1}{3} e^{-t+\tau} \\ \frac{2}{3} e^{5t-5\tau} - \frac{2}{3} e^{-t+\tau} & \frac{2}{3} e^{5t-5\tau} + \frac{1}{3} e^{-t+\tau} \end{bmatrix} \quad (19)$$

>  $ProdTau := evalm(MatExpTau &* BBtau) : ProdTau[1, 1]; ProdTau[2, 1]$

$$\begin{aligned} 5 \left( \frac{1}{3} e^{5t-5\tau} + \frac{2}{3} e^{-t+\tau} \right) e^{3\tau} - 7 \left( \frac{1}{3} e^{5t-5\tau} - \frac{1}{3} e^{-t+\tau} \right) e^{2\tau} \\ 5 \left( \frac{2}{3} e^{5t-5\tau} - \frac{2}{3} e^{-t+\tau} \right) e^{3\tau} - 7 \left( \frac{2}{3} e^{5t-5\tau} + \frac{1}{3} e^{-t+\tau} \right) e^{2\tau} \end{aligned} \quad (20)$$

>  $MatExpNoHom := map(int, ProdTau, tau=0..t)$

$$MatExpNoHom := \begin{bmatrix} \frac{1}{18} (e^{6t} + 28 e^{3t} - 29) e^{-t} \\ -\frac{1}{18} (-2 e^{6t} + 45 e^{4t} - 14 e^{3t} - 29) e^{-t} \end{bmatrix} \quad (21)$$

>  $Compr := map(rcurry(eval, t=0'), MatExpNoHom)$

$$Compr := \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad (22)$$

>  $Sol[1] := x[1](t) = simplify(MatExpHom[1, 1] + MatExpNoHom[1, 1])$

$$Sol_1 := x_1(t) = \frac{7}{18} e^{-t} + \frac{1}{18} e^{5t} + \frac{14}{9} e^{2t} \quad (23)$$

>  $Sol[2] := x[2](t) = simplify(MatExpHom[2, 1] + MatExpNoHom[2, 1])$

$$Sol_2 := x_2(t) = -\frac{7}{18} e^{-t} + \frac{1}{9} e^{5t} - \frac{5}{2} e^{3t} + \frac{7}{9} e^{2t} \quad (24)$$

>  $Sist[1]$

$$\frac{d}{dt} x_1(t) = x_1(t) + 2 x_2(t) + 5 e^{3t} \quad (25)$$

>  $Sist[2]$

$$\frac{d}{dt} x_2(t) = 4 x_1(t) + 3 x_2(t) - 7 e^{2t} \quad (26)$$

>  $CompUno := simplify(eval(subs(x[1](t) = rhs(Sol[1]), x[2](t) = rhs(Sol[2]), lhs(Sist[1]) - rhs(Sist[1]) = 0)))$

$$CompUno := 0 = 0 \quad (27)$$

>  $CompDos := simplify(eval(subs(x[1](t) = rhs(Sol[1]), x[2](t) = rhs(Sol[2]), lhs(Sist[2]) - rhs(Sist[2]) = 0)))$

$$CompDos := 0 = 0 \quad (28)$$

>  $CompTres := simplify(subs(t=0, Sol[1]))$

$$CompTres := x_1(0) = 2 \quad (29)$$

| > *CompCuatro* := *simplify*(*subs*(*t*=0, *Sol*[2]))  
| > *CompCuatro* := *x*<sub>2</sub>(0) = -2  
| >

(30)